



What to Expect with EPKINLY

A guide to treatment



Actor portrayals.

What is EPKINLY?

EPKINLY is a prescription medicine used to treat adults with certain types of diffuse large B-cell lymphoma (DLBCL), high-grade B-cell lymphoma, or follicular lymphoma (FL) that has come back or that did not respond to previous treatment after receiving 2 or more treatments. EPKINLY is approved based on patient response data. Studies are ongoing to confirm the clinical benefit of EPKINLY. It is not known if EPKINLY is safe and effective in children.

SELECT IMPORTANT SAFETY INFORMATION

EPKINLY can cause serious side effects, including:

- Cytokine release syndrome (CRS), which is common during treatment with EPKINLY and can be serious or life-threatening.
- Neurologic problems that can be life-threatening and lead to death.
- Infections and Low blood cell counts.

What's inside this guide?

Welcome to EPKINLY
About EPKINLY
What to do before treatment
Important Safety Information
What to expect during treatment
What to do after treatment
Resources
Treatment Tracker
Appointment Calendar 24



Care Partner tip: Read this guide with your loved one so that you can plan ahead together and get the most out of their treatment with EPKINLY.





Welcome to EPKINLY

Starting a new treatment can prompt a lot of questions. This guide is designed to help you and your Care Partner know what to expect before, during, and after EPKINLY treatment.

- Use the tabs, or the table of contents, to easily find what you're looking for
- Use the page in the back of the binder to keep track of your treatment team
- Remember to download and print two wallet cards (one for you and one for your Care Partner)



Scan the QR code or visit <u>EPKINLY.com</u> for more information.







About EPKINLY



A bispecific antibody treatment that harnesses the power of your own immune system to help fight cancer



Given as an injection, it's not an infusion, chemotherapy, or stem cell therapy



A treatment that may be given at a location that is able to administer EPKINLY, such as your oncologist's office or a nearby outpatient center

SELECT IMPORTANT SAFETY INFORMATION


Important Warnings—EPKINLY can cause serious side effects, including:

- Cytokine release syndrome (CRS), which is common during treatment with EPKINLY and can be serious or life-threatening. To help reduce your risk of CRS, you will receive EPKINLY on a step-up dosing schedule (when you receive 2 or 3 smaller step-up doses of EPKINLY before your first full dose during your first cycle of treatment), and you may also receive other medicines before and for 3 days after receiving EPKINLY. If your dose of EPKINLY is delayed for any reason, you may need to repeat the step-up dosing schedule.
- **Neurologic problems** that can be life-threatening and lead to death. Neurologic problems may happen days or weeks after you receive EPKINLY.



How often you will receive EPKINLY

The following outlines a general schedule for when you will receive your treatment. Work with your healthcare provider to determine your appointment dates. Your EPKINLY treatment schedule is divided into cycles that are usually 28 days (4 weeks) long.



- In the beginning, you will receive EPKINLY based on a "step-up dosing schedule." The step-up dosing schedule is when you receive 2 or 3 smaller "step-up" doses of EPKINLY during your first cycle of treatment. Step-up dosing is designed to help you adjust to treatment safely. This helps reduce the possibility of a certain side effect called cytokine release syndrome (CRS)
- Before and for 3 days after each dose of EPKINLY in your first 4 weeks of treatment, you will receive other medicines to help reduce your risk of CRS. Your healthcare provider will decide if you need to receive medicine to help reduce your risk of CRS in later doses of EPKINLY
- You may have side effects during treatment. Your healthcare provider will monitor you for symptoms of CRS, neurologic problems, infections, and low blood cell counts. Your treatment may be temporarily or completely stopped if you develop certain side effects
- Staying on treatment with EPKINLY may help you achieve remission and continue to see response

SELECT IMPORTANT SAFETY INFORMATION

People with DLBCL or high-grade B-cell lymphoma should be hospitalized for 24 hours after receiving their first full dose of EPKINLY on day 15 of cycle 1 due to the risk of CRS and neurologic problems.



What to do before treatment

There are some helpful things to know before your visit. Your healthcare provider may also have additional instructions.



Hydration

It is important that you are adequately hydrated before you receive your EPKINLY injection.

Start drinking plenty of water the day before your visit. Consider bringing water with you to your visit.



Injection site

Your EPKINLY injection will likely be in the lower abdomen or the thigh.

- Wear loose clothing that is easy to change
- Prepare a cold pack (included in this binder) in your freezer in case of any injection site reaction after treatment

For Patients with Diffuse Large B-cell Lymphoma (DLBCL) or High-grade B-cell Lymphoma



Planning for your first full dose

Due to the risk of cytokine release syndrome (CRS) and neurologic problems, you should be hospitalized for 24 hours after receiving your first full dose of EPKINLY in week 3.

 Make sure you discuss with your healthcare provider where you will need to receive this dose so you can plan ahead if this is not near where you live



Your healthcare provider may receive EPKINLY from a specialty pharmacy, depending on your health insurance. Ask your healthcare provider about how he or she will receive your treatment. You may get a call from the pharmacy via a 1-800 number to collect any co-pay or co-insurance.

SELECT IMPORTANT SAFETY INFORMATION

Tell your healthcare provider or get medical help right away if you develop a fever of 100.4°F (38°C) or higher; dizziness or lightheadedness; trouble breathing; chills; fast heartbeat; feeling anxious; headache; confusion; shaking (tremors); problems with balance and movement, such as trouble walking; trouble speaking or writing; confusion and disorientation; drowsiness, tiredness or lack of energy; muscle weakness; seizures; or memory loss. **These may be symptoms of CRS or neurologic problems.** If you have any symptoms that impair consciousness, **do not** drive or use heavy machinery or do other dangerous activities until your symptoms go away.

Plan for your treatment day

What should I ask?

Before your appointment for your treatment with EPKINLY, ask your healthcare provider questions to help you plan and prepare.

What do I need to prepare for my visit?
Where should I go for my treatment?
Will I need my Care Partner to drive me to and from my injection visits?
How long will my visit last? Will I be able to go home right after my treatment?
Notes:



Important Safety Information

What is EPKINLY?

EPKINLY is a prescription medicine used to treat adults with certain types of diffuse large B-cell lymphoma (DLBCL), high-grade B-cell lymphoma, or follicular lymphoma (FL) that has come back or that did not respond to previous treatment after receiving 2 or more treatments. EPKINLY is approved based on patient response data. Studies are ongoing to confirm the clinical benefit of EPKINLY. It is not known if EPKINLY is safe and effective in children.

IMPORTANT SAFETY INFORMATION Important Warnings—EPKINLY can cause serious side effects, including:

- Cytokine release syndrome (CRS), which is common during treatment with EPKINLY and can be serious or life-threatening. To help reduce your risk of CRS, you will receive EPKINLY on a step-up dosing schedule (when you receive 2 or 3 smaller step-up doses of EPKINLY before your first full dose during your first cycle of treatment), and you may also receive other medicines before and for 3 days after receiving EPKINLY. If your dose of EPKINLY is delayed for any reason, you may need to repeat the step-up dosing schedule.
- Neurologic problems that can be life-threatening and lead to death.
 Neurologic problems may happen days or weeks after you receive EPKINLY.

People with DLBCL or high-grade B-cell lymphoma should be hospitalized for 24 hours after receiving their first full dose of EPKINLY on day 15 of cycle 1 due to the risk of CRS and neurologic problems.

Tell your healthcare provider or get medical help right away if you develop a fever of 100.4°F (38°C) or higher; dizziness or lightheadedness; trouble breathing; chills; fast heartbeat; feeling anxious; headache; confusion; shaking (tremors); problems with balance and movement, such as trouble walking; trouble speaking or writing; confusion and disorientation; drowsiness, tiredness or lack of energy; muscle weakness; seizures; or memory loss. These may be symptoms of CRS or neurologic

problems. If you have any symptoms that impair consciousness, **do not** drive or use heavy machinery or do other dangerous activities until your symptoms go away.

EPKINLY can cause other serious side effects, including:

- Infections that may lead to death. Your healthcare provider will check you for signs and symptoms of infection before and during treatment and treat you as needed if you develop an infection. You should receive medicines from your healthcare provider before you start treatment to help prevent infection. Tell your healthcare provider right away if you develop any symptoms of infection during treatment, including fever of 100.4°F (38°C) or higher, cough, chest pain, tiredness, shortness of breath, painful rash, sore throat, pain during urination, or feeling weak or generally unwell.
- Low blood cell counts, which can be serious or severe. Your healthcare provider will check your blood cell counts during treatment. EPKINLY may cause low blood cell counts, including low white blood cells (neutropenia), which can increase your risk for infection; low red blood cells (anemia), which can cause tiredness and shortness of breath; and low platelets (thrombocytopenia), which can cause bruising or bleeding problems.

Your healthcare provider will monitor you for symptoms of CRS, neurologic problems, infections, and low blood cell counts during treatment with EPKINLY. Your healthcare provider may temporarily stop or completely stop treatment with EPKINLY if you develop certain side effects.



Important Safety Information (continued)


Before you receive EPKINLY, tell your healthcare provider about all your medical conditions, including if you have an infection, are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed. If you receive EPKINLY while pregnant, it may harm your unborn baby. If you are a female who can become pregnant, your healthcare provider should do a pregnancy test before you start treatment with EPKINLY and you should use effective birth control (contraception) during treatment and for 4 months after your last dose of EPKINLY. Tell your healthcare provider if you become pregnant or think that you may be pregnant during treatment with EPKINLY. Do not breastfeed during treatment with EPKINLY and for 4 months after your last dose of EPKINLY.

In DLBCL or high-grade B-cell lymphoma, the most common side effects of EPKINLY include CRS, tiredness, muscle and bone pain, injection site reactions, fever, stomach-area (abdominal) pain, nausea, and diarrhea. The most common severe abnormal laboratory test results include decreased white blood cells, decreased red blood cells, and decreased platelets.

In follicular lymphoma the most common side effects of EPKINLY include injection site reactions, CRS, COVID-19, tiredness, upper respiratory tract infections, muscle and bone pain, rash, diarrhea, fever, cough, and headache. The most common severe abnormal laboratory test results include decreased white blood cells and decreased red blood cells.

These are not all of the possible side effects of EPKINLY. Call your doctor for medical advice about side effects. You are encouraged to report side effects to the FDA at (800) FDA-1088 or www.fda.gov/medwatch or to Genmab US, Inc. at 1-855-4GENMAB (1-855-443-6622).

Please see <u>Medication Guide</u>, including Important Warnings.







About your injection

Before your healthcare provider administers your injection, he or she will:


- Check your vital signs (temperature, blood pressure, heart rate) and ask you some questions about your general health status
- Check to make sure you are adequately hydrated
- Before each dose of EPKINLY in the first 4 weeks, you will receive medications 30 to 120 minutes before your injection to help reduce your risk of CRS and you will also receive medication for 3 days after each dose. Your healthcare provider will decide if you need to receive medicine to help reduce your risk of CRS with future doses

During your injection

- EPKINLY will be given as an injection under your skin (subcutaneous injection), usually in the lower part of your stomach area (abdomen) or thigh
- Your healthcare provider will likely switch the injection site at each visit (left to right side or vice versa), especially during your first few weeks of treatment
- You may be asked by your healthcare team to stay longer after the injection so that they can monitor for any reactions

Keep track of your injection site locations by using the Treatment Tracker on page 18.







Care Partner tip:

- A cold pack is included in this binder to help with any injection site reactions related to treatment. Remember to prepare the cold pack in the freezer ahead of time
- Make sure your transportation arrangements can accommodate staying longer after the injection if the healthcare team requests it

SELECT IMPORTANT SAFETY INFORMATION

EPKINLY can cause serious side effects, including:

- Cytokine release syndrome (CRS), which is common during treatment with EPKINLY and can be serious or life-threatening.
- **Neurologic problems** that can be life-threatening and lead to death.
- Infections and Low blood cell counts.





Approved for adults only.

Questions and notes

Following up with your healthcare team

After your first treatment visit for EPKINLY, consider discussing the following with your healthcare team:

What should I do if I miss a dose?
Is there anything I need to be aware of while I am receiving EPKINLY?
Is there anything I can do to make my overall health better during treatment?



Questions and notes

Write down questions you may have before or during your appointment with your healthcare provider. You can also use this space to take notes.						



What to do in between treatments

After you receive a dose of EPKINLY, your healthcare provider will monitor you for any side effects from the treatment, and you should also look out for side effects and any changes in how you're feeling afterward.

EPKINLY can cause serious side effects, including cytokine release syndrome (called CRS), which is common during EPKINLY treatment and can be serious or life-threatening, and neurologic problems that can be life-threatening and lead to death. Other serious side effects include infections that can be serious and lead to death and low blood cell counts, which can be serious or severe.

Watch for symptoms of CRS

CRS symptoms may include

- Fever 100.4°F (38°C) or higher
- Dizziness or lightheadedness
- Trouble breathing
- Chills
- Fast heartbeat
- Feeling anxious

- Headache
- Confusion
- Shaking (tremors)
- Problems with balance and movement, such as trouble walking

Tell your healthcare provider or get medical help right away if you develop symptoms of CRS. Your healthcare provider may give you medicines to treat your side effects. Your healthcare provider may temporarily or completely stop your treatment with EPKINLY, depending on your symptoms.



Care Partner tip: Monitor your loved one for signs of fever. Use a thermometer to check frequently for fever of 100.4°F (38°C) or higher in the days following an injection of EPKINLY. Alert the healthcare team if your loved one has a fever of 100.4°F (38°C) or higher following treatment.

Download two copies of your **wallet card** (one for you and one for your Care Partner) and carry them with you in the event a healthcare provider needs quick access to your information.

SELECT IMPORTANT SAFETY INFORMATION

In DLBCL or high-grade B-cell lymphoma, the most common side effects of EPKINLY include CRS, tiredness, muscle and bone pain, injection site reactions, fever, stomach-area (abdominal) pain, nausea, and diarrhea. The most common severe abnormal laboratory test results include decreased white blood cells, decreased red blood cells, and decreased platelets.

In follicular lymphoma the most common side effects of EPKINLY include injection site reactions, CRS, COVID-19, tiredness, upper respiratory tract infections, muscle and bone pain, rash, diarrhea, fever, cough, and headache. The most common severe abnormal laboratory test results include decreased white blood cells and decreased red blood cells.

Watch for symptoms of neurologic problems

Neurologic problems may happen days or weeks after you receive EPKINLY. Your healthcare provider may refer you to a healthcare provider who specializes in neurologic problems.

Symptoms may include but are not limited to:

- Trouble speaking or writing
- Confusion and disorientation
 Shaking (tremors)
- Drowsiness

- Muscle weakness
- Seizures
- Tiredness or lack of energy
- Memory loss

Tell your healthcare provider right away if you develop any symptoms of neurologic problems. You may be given medicines to treat your side effects, and your treatment with EPKINLY may be temporarily or completely stopped.



Care Partner tip: Neurologic problems can affect your loved one's ability to think, remember, and/or use language (verbal or written). These may be subtle or hard to notice at first. If you notice any changes in behavior in your loved one, let the healthcare team know immediately.

Additional important things to know

People with diffuse large B-cell lymphoma (DLBCL) or high-grade B-cell lymphoma should be hospitalized for 24 hours after receiving their first full dose of EPKINLY on Day 15 of Cycle 1 due to the risk of CRS and neurologic problems.

Do not drive, operate heavy machinery, or do other dangerous activities if you develop dizziness, confusion, tremors, sleepiness, or any other symptoms that impair consciousness until your signs and symptoms go away. These may be signs and symptoms of CRS or neurologic problems.

Watch for symptoms of infections. Your healthcare provider will check you for signs and symptoms of infection before and during treatment and treat you as needed if you develop an infection. You should receive medicines from your healthcare provider before you start treatment to help prevent infection.

Symptoms may include but are not limited to: fever of 100.4°F (38°C) or higher; cough; chest pain; tiredness; shortness of breath; painful rash; sore throat; pain during urination; feeling weak or generally unwell.

Tell your healthcare provider right away if you develop symptoms of an infection. Your healthcare provider may temporarily or completely stop your treatment with EPKINLY. These are not all the possible side effects of EPKINLY. Call your doctor for medical advice about side effects.



MyNavCare[™] is here to support you



With *MyNavCare*, you can receive personalized support throughout your treatment journey from a dedicated team that cares



A partner on your treatment journey

Your healthcare provider is your main source for treatment information. With *MyNavCare*, you gain access to additional resources and medication information.

- Dedicated support throughout your treatment
- Information about treatment access and financial support*
- Insight into independent organizations that may provide additional assistance
- Additional tools and resources that may help along the way



Financial assistance for eligible patients

MyNavCare may be able to help uninsured or underinsured patients receive financial assistance and information. This includes:

- Co-pay assistance program for commercially insured patients*
- Independent patient assistant foundation information[†]
- Genmab Patient Assistance Program



Ongoing personalized support

Your **Patient Engagement Liaison**[†] provides ongoing support throughout treatment. They're dedicated to you, helping to:

- Provide information about your condition
- Connect you to third-party organizations that may be able to help
- Offer resources for you and your Care Partners

Contact a *MyNavCare* Support Specialist by calling 1-866-NAV-CAR1 (1-866-628-2271), Monday-Friday, 8 AM - 8 PM ET. You can also scan the QR code or visit <u>www.MyNavCare.com</u> to learn more.





^{*}Based on eligibility requirements and Terms and Conditions.

[†]Eligibility criteria are determined by each independent third-party foundation. *MyNavCare* does not determine eligibility and cannot guarantee foundations will offer assistance.

^{*}MyNavCare Patient Engagement Liaisons are part of the MyNavCare Patient Support Program and do not provide medical advice or work under the direction of the prescribing healthcare providers. They are trained to direct patients to speak with their healthcare provider about any treatment-related questions.

Additional resources

Support for you and your Care Partner

Here are organizations that can help patients and Care Partners connect with others, and provide information and resources throughout treatment.





Cancer Care

www.cancercare.org | 800-813-HOPE (4673)



American Cancer Society

www.cancer.org | 800-227-2345



Lymphoma Research Foundation

www.lymphoma.org | 800-500-9976



The Leukemia & Lymphoma Society

www.lls.org | 800-955-4572



Family Reach

www.familyreach.org | 973-394-1411 ext. 1



Follicular Lymphoma Foundation

www.theflf.org | info@theflf.org

There are many available resources that may be helpful to patients, families, and Care Partners dealing with cancer. The above resources are not controlled, owned, or endorsed by Genmab/AbbVie, and Genmab/AbbVie is not responsible for their content.





If you are a Care Partner

Some responsibilities you may take on during your loved one's treatment:



Decision-making: Researching clinical trials and treatment options, helping to consider treatment choices, and being an advocate for your loved one



Emotional: Encouraging your loved one to share their feelings and helping them feel supported



Medical: Helping watch for potential side effects, attending appointments, and ensuring they have what they need throughout their treatments



Logistical: Scheduling and driving them to appointments, running errands, and assisting with daily needs





Track important information here

Record your experience with EPKINLY and share with your healthcare team so they get a full picture of how you're doing on treatment.

- Keep track of the appointment date for each dose
- Mark the location of the injection site
- Write down how you're feeling after the injection and any other information you want to share

DOSE	How I'm feeling:
Date:	
Time:	
Injection Site:	
	Questions/notes for my healthcare team:
	<u></u>
// \	
DOSE	How I'm feeling:
Date:	
Time:	
Injection Site:	
	Questions/notes for my healthcare team:
	Questions/hotes for my heatthcare team.
//_//	
U = U	
DOSE	How I'm feeling:
Date:	
Time:	
Injection Site:	
	Questions/notes for my healthcare team:
// /	
(/\=\\	





DOSE	How I'm feeling:
Date:	
Time:	
Injection Site:	
	Questions/notes for my healthcare team:
$\sqrt{\Lambda}$	
POSE	How the feelings
Dose	How I'm feeling:
Time:	
Injection Site:	
	Questions/notes for my healthcare team:
\int_{Λ}	
DOSE	How I'm feeling:
Date:	
Time:	
Injection Site:	
	Questions/notes for my healthcare team:
/ \ •\\	





DOSE	How I'm feeling:
Date:	
Time:	
Injection Site:	
	Questions/notes for my healthcare team:
DOSE	How I'm feeling:
Date:	
Time:	
Injection Site:	
	Questions/notes for my healthcare team:
/ / •/\	
131131	
DOSE	How I'm feeling:
Date:	-
Time:	
Injection Site:	
	Questions/notes for my healthcare team:
///\	
UTIV	





DOSE	How I'm feeling:
Date:	
Time:	
Injection Site:	
	Questions/notes for my healthcare team:
DOSE	How I'm feeling:
Date:	
Time:	
Injection Site:	
	Questions/notes for my healthcare team:
/ / •/\	
131131	
DOSE	How I'm feeling:
Date:	
Time:	
Injection Site:	
	Questions/notes for my healthcare team:
///\	
$U \prod V$	





DOSE	How I'm feeling:
Date:	
Time:	
Injection Site:	
	Questions/notes for my healthcare team:
DOSE	How I'm feeling:
Date:	
Time:	
Injection Site:	
	Questions/notes for my healthcare team:
// <u>~</u> //	
	the Berteeling
DOSE	How I'm feeling:
Date:	
Time:	
Injection Site:	
	Questions/notes for my healthcare team:
// //	
101101	





DOSE	How I'm feeling:
Date:	
Time:	
Injection Site:	
	Questions/notes for my healthcare team:
DOSE	How I'm feeling:
Date:	
Time:	
Injection Site:	
	Questions/notes for my healthcare team:
/ / •/\	
131131	
DOSE	How I'm feeling:
Date:	
Time:	
Injection Site:	
	Questions/notes for my healthcare team:
///\	
$U \prod V$	





Calendar

Use this calendar to keep track of your healthcare appointments, including injections and tests. Make notes here prior to the appointment if you have questions for your healthcare provider.

MONTH:								
SUN	MON	TUE	WED	THU	FRI	SAT		
Notes								





MONTH:									
SUI	N	MON	TUE	WED	THU	FRI	SAT		

Notes		





MONTI	MONTH:										
SUN	MON	TUE	WED	THU	FRI	SAT					

Notes			





			_										
N	MONTH:												
SUN	N	МО	N	TUE		WEI	D	THU	J	FRI		SAT	

Notes			





MONTI	MONTH:									
SUN	MON	TUE	WED	THU	FRI	SAT				
Notes										





MONTI	MONTH:										
SUN	MON	TUE	WED	THU	FRI	SAT					
Notes											

epkinly[®] epcoritamab-bysp

SAT



MONTI	H:							
SUN	MON	TUE	WEI	D	THU	J	FRI	

				•		
				'		

Notes			





MONTI	Н:					
SUN	MON	TUE	WED	THU	FRI	SAT
Notes						

epkinly[®] epcoritamab-bysp



MONTI	н:				
SUN	MON	TUE	WED	THU	FRI

SUN	MON	TUE	WED	THU	FRI	SAT

Notes			





MONTI	Н:				
SUN	MON	TUE	WED	THU	FRI

SUN	MON	TUE	WED	THU	FRI	SAT

Notes			





MONTI	Н:					
SUN	MON	TUE	WED	THU	FRI	SAT
Notos	1		1			1

NOTES		





MC	ONTH:					
SUN	MON	TUE	WED	THU	FRI	SAT

Notes		





M	IONTI	H:					
SUN		MON	TUE	WED	THU	FRI	SAT

Notes			



Questions and notes

Write down questions you may have before or during your appointment with your healthcare provider. You can also use this space to take notes.



Important contact information

Write down contact information for members of your healthcare team and other emergency contacts for quick and easy access.

My Doctor
Name:
Phone number:
Office address:
My Nurses
Name:
Phone number:
Office address:
Name:
Phone number:
Office address:
NA v Dis cryps cross
My Pharmacy
Name:
Phone number:
Pharmacy address:
My Emergency Contact
Name:
Phone number:
If you notice any side effects or unusual symptoms, reach out to your healthcare team right away.



It's a good idea to keep this info in your mobile phone as well. (You can even just take a picture of this page.)







Approved for adults only.



Scan the QR code or visit <u>EPKINLY.com</u> for more information.



